



ACC.14

TCT@ACC-i2 | innovation in intervention

A1690

JACC April 1, 2014

Volume 63, Issue 12



TCT@ACC-i2: The Interventional Learning Pathway

LONG-TERM OUTCOME AFTER DEFERRAL OF REVASCULARIZATION IN PATIENTS WITH INTERMEDIATE CORONARY STENOSIS AND GRAY-ZONE FRACTIONAL FLOW RESERVE

Oral Contributions

Room 209 C

Saturday, March 29, 2014, 8:30 a.m.-8:40 a.m.

Session Title: Intravascular Physiology and Imaging

Abstract Category: 35. TCT@ACC-i2: IVUS and Intravascular Physiology

Presentation Number: 2902-05

Authors: *Yasutsugu Shiono, Takashi Kubo, Yousuke Katayama, Tomoko Tohoda, Yuki Nozawa, Mao Yokoyama, Yuichi Ozaki, Makoto Orii, Kunihiro Shimamura, Takashi Yamano, Takashi Tanimoto, Yoshiki Matsuo, Yasushi Ino, Tomoyuki Yamaguchi, Kumiko Hirata, Atsushi Tanaka, Toshio Imanishi, Takashi Akasaka, Wakayama Medical University, Wakayama, Japan*

Background: The safety of deferring revascularization of intermediate coronary stenosis with fractional flow reserve (FFR) of 0.75 - 0.80, the so-called "gray zone", remains debatable. The aim of this study was to assess the safety of deferring revascularization for patients with FFR of 0.75 - 0.80 compared with those with FFR of >0.80.

Methods: We assessed long-term outcomes of 150 patients with angiographically intermediate stenosis deferred from revascularization on the basis of FFR ≥ 0.75 . Target vessel failures (TVF) defined as a composite of cardiac death, target vessel related myocardial infarction, and ischemia-driven target vessel revascularization were evaluated during follow-up period.

Results: A total of 56 patients had coronary lesions with FFR of 0.75 - 0.80 and 94 patients had those with FFR of >0.80. There was no difference in baseline clinical characteristics between the two groups. Patients with FFR of 0.75 - 0.80 had more left anterior descending lesions than those with FFR >0.80 (75% vs. 44%, $p = 0.001$). During a median follow-up period of 3.0 (2.1 - 4.0) years, the incidence of TVF was higher in patients with FFR of 0.75 - 0.80 than those with FFR of >0.80 (16% vs. 3%; hazard ratio, 5.2; 95% confidence interval, 1.4 to 19.5; $p = 0.015$). All TVF consisted of target vessel revascularization, except for one case of cardiac death in patients with FFR of 0.75 - 0.80.

Conclusions: Patients with FFR of 0.75 - 0.80 were at higher risk of TVF mainly due to target vessel revascularization than those with FFR of >0.80.